



Model created in COMSOL Multiphysics 6.4

# Ion Cyclotron Motion

## Introduction

---

If a charged particle is placed in a uniform magnetic field, it moves in a helical pattern about a fixed gyro radius. The gyro radius, which is also known as the Larmor or cyclotron radius, is given by the simple equation:

$$r_L = \frac{mv_{\perp}}{ZeB}$$

- $r_L$  (SI unit: m) is the Larmor radius,
- $v_{\perp}$  (SI unit: m/s) is the velocity component orthogonal to the magnetic field,
- $Z$  (dimensionless) is the particle charge number,
- $e = 1.602176634 \times 10^{-19}$  C is the elementary charge,
- $m$  (SI unit: kg) is the particle mass, and
- $B$  (SI unit: T) is the magnitude of the magnetic flux density.

This model computes the trajectory of an ion in a uniform magnetic field using the Newtonian, Lagrangian, and Hamiltonian formulations available in the Mathematical Particle Tracing interface.

## Model Definition

---

The equations of motion for a charge in a magnetic field can be determined from the Lagrange equations:

$$\frac{d}{dt} \left( \frac{\partial L}{\partial \mathbf{v}} \right) = \frac{\partial L}{\partial \mathbf{q}} \quad (1)$$

where  $\mathbf{v}$  is the particle velocity,  $\mathbf{q}$  is the particle position, and  $L$  (SI unit: J) is the Lagrangian, which is defined as:

$$L = \frac{m(\mathbf{v} \cdot \mathbf{v})}{2} + q(\mathbf{v} \cdot \mathbf{A})$$

This form of the Lagrangian is valid for nonrelativistic particles; that is, the particle velocity is much less than the speed of light. The contribution due to the electric potential is neglected. The Hamiltonian is related to the Lagrangian via:

$$H = \mathbf{v} \cdot \frac{\partial L}{\partial \mathbf{v}} - L$$

Introducing the generalized momentum of the particle,  $\mathbf{P}$  (SI unit: kg·m/s), the Hamiltonian becomes:

$$H = \frac{(\mathbf{P} - q\mathbf{A})^2}{2m}$$

In order to derive the equations of motion for the Newtonian formulation, start with the right-hand side of [Equation 1](#):

$$\frac{\partial L}{\partial \mathbf{q}} = \nabla L = q\nabla(\mathbf{A} \cdot \mathbf{v}) = q(\mathbf{v} \cdot \nabla)\mathbf{A} + q(\mathbf{v} \times \nabla \times \mathbf{A}) \quad (2)$$

The left-hand side of [Equation 1](#) becomes

$$\frac{d}{dt} \left( \frac{\partial L}{\partial \mathbf{v}} \right) = \frac{d}{dt} (\mathbf{p} + q\mathbf{A}) = \frac{d\mathbf{p}}{dt} + q(\mathbf{v} \cdot \nabla)\mathbf{A} + q\frac{\partial \mathbf{A}}{\partial t} . \quad (3)$$

Equating [Equation 2](#) and [Equation 3](#) and canceling like terms yields

$$\frac{d\mathbf{p}}{dt} = q(\mathbf{v} \times \mathbf{B}) \quad (4)$$

for a stationary magnetic field. Here, the magnetic flux density has been introduced as

$$\mathbf{B} = \nabla \times \mathbf{A} \quad (5)$$

When the particle velocity is small compared to the speed of light [Equation 4](#) yields the classical equation of motion for a charged particle in a stationary, uniform magnetic field

$$\frac{d}{dt}(m\mathbf{v}) = q(\mathbf{v} \times \mathbf{B})$$

## *Results and Discussion*

The model is solved in COMSOL using the Lagrangian, Hamiltonian, and Newtonian formulations. The Larmor radius is compared to the analytic solution and given in [Table 1](#). All formulations agree with the analytic expression to within 0.05%. The two coupled first-order differential equations give a slightly different result from a single second-order differential equation for each coordinate.

TABLE 1: TABLE COMPARING THE LARMOR RADIUS FOR THE DIFFERENT FORMULATIONS.

	ANALYTIC	LAGRANGIAN	HAMILTONIAN	NEWTONIAN, 2ND ORDER	NEWTONIAN, 1ST ORDER
Larmor radius ( $\mu\text{m}$ )	414.57	414.55	414.55	414.55	414.57

The particle trajectories for the three different formulations are plotted below.




Figure 1: Plot of the ion trajectory for the Lagrangian formulation.




Figure 2: Plot of the ion trajectory for the Hamiltonian formulation.




Figure 3: Plot of the particle trajectory for the Newtonian formulation.




Figure 4: Plot of the particle trajectory for the first-order Newtonian formulation.

## Reference

---

1. L.D. Landau and E.M. Lifshitz, *The Classical Theory of Fields*, 4th ed., Elsevier, 2005.

---

**Application Library path:** Particle\_Tracing\_Module/  
Charged\_Particle\_Tracing/ion\_cyclotron\_motion


---

## Modeling Instructions




---

From the **File** menu, choose **New**.

### NEW



In the **New** window, click  **Model Wizard**.

### MODEL WIZARD

- 1 In the **Model Wizard** window, click  **3D**.
- 2 In the **Select Physics** tree, select **Mathematics > Mathematical Particle Tracing (pt)**.
- 3 Click **Add**.
- 4 Click  **Study**.
- 5 In the **Select Study** tree, select **General Studies > Time Dependent**.
- 6 Click  **Done**.

### GEOMETRY 1

*Cylinder 1 (cyl1)*

- 1 In the **Geometry** toolbar, click  **Cylinder**.
- 2 In the **Settings** window for **Cylinder**, locate the **Size and Shape** section.
- 3 In the **Radius** text field, type  $2e-3$ .
- 4 In the **Height** text field, type  $2e-3$ .
- 5 In the **Geometry** toolbar, click  **Build All**.

### GLOBAL DEFINITIONS

Define parameters for the particle mass, magnetic flux density, initial particle velocity, and Larmor radius. The Larmor radius is only used during results processing.

*Parameters 1*

- 1 In the **Model Builder** window, under **Global Definitions** click **Parameters 1**.
- 2 In the **Settings** window for **Parameters**, locate the **Parameters** section.
- 3 In the table, enter the following settings:

Name	Expression	Value	Description
mp	$0.04[\text{kg/mol}]/N\_A\_const$	6.6422E-26 kg	Ion mass
B	$2[\text{T}]$	2 T	Magnetic flux density
v0	$2E3[\text{m/s}]$	2000 m/s	Particle velocity, perpendicular to the magnetic field
rL	$mp*v0/(e\_const*B)$	4.1457E-4 m	Larmor radius

**DEFINITIONS**

Define an analytic expression for the magnetic vector potential, which results in a uniform magnetic field in the  $z$  direction.

*Variables 1*


- 1 In the **Model Builder** window, expand the **Component 1 (comp1) > Definitions** node.
- 2 Right-click **Definitions** and choose **Local Variables**.
- 3 In the **Settings** window for **Variables**, locate the **Variables** section.
- 4 In the table, enter the following settings:

Name	Expression	Unit	Description
Ax	$1[\text{Wb/m}]*y[1/\text{m}]$	Wb/m	Magnetic vector potential, x-component
Ay	$-1[\text{Wb/m}]*x[1/\text{m}]$	Wb/m	Magnetic vector potential, y-component
Az	$0[\text{Wb/m}]$	Wb/m	Magnetic vector potential, z-component
Bx	$d(Az, y) - d(Ay, z)$	T	Magnetic flux density, x-component
By	$d(Ax, z) - d(Az, x)$	T	Magnetic flux density, y-component
Bz	$d(Ay, x) - d(Ax, y)$	T	Magnetic flux density, z-component

## MATHEMATICAL PARTICLE TRACING (PT)

Release a single particle at the origin with an initial velocity in the  $x$  direction so that the Lorentz force is nonzero. Also add a small initial velocity in the  $z$  direction so that you can clearly see the particle trajectory after solving.

### *Release from Grid 1*

- 1 In the **Physics** toolbar, click  **Global** and choose **Release from Grid**.
- 2 In the **Settings** window for **Release from Grid**, locate the **Initial Velocity** section.
- 3 Specify the  $\mathbf{v}_0$  vector as

$v_0$	$x$
0	$y$
$1e2$	$z$

The first formulation you will use is **Lagrangian**. The Lagrangian for a particle in a magnetic field is the sum of the particle kinetic energy, which is here defined as  $\mathbf{pt} \cdot \mathbf{Ep}$ , and the dot product of the particle velocity and the magnetic potential, multiplied by the particle charge.


- 4 In the **Model Builder** window, click **Mathematical Particle Tracing (pt)**.
- 5 In the **Settings** window for **Mathematical Particle Tracing**, locate the **Particle Release and Propagation** section.
- 6 From the **Formulation** list, choose **Lagrangian**.

### *Particle Properties 1*

- 1 In the **Model Builder** window, click **Particle Properties 1**.
- 2 In the **Settings** window for **Particle Properties**, locate the **Particle Mass** section.
- 3 In the  $m_p$  text field, type `mp`.
- 4 Locate the **Lagrangian** section. In the  $L$  text field, type `pt.Ep+e_const*(pt.vx*Ax+pt.vy*Ay+pt.vz*Az)`.


## MESH 1

Use a coarse mesh. The field is entered using an analytic expression, so the accuracy of the solution is independent of the mesh element size.

- 1 In the **Model Builder** window, under **Component 1 (comp1)** click **Mesh 1**.
- 2 In the **Settings** window for **Mesh**, locate the **Physics-Controlled Mesh** section.
- 3 From the **Element size** list, choose **Extra coarse**.
- 4 Click  **Build All**.

## STUDY 1

### *Step 1: Time Dependent*

- 1 In the **Model Builder** window, under **Study 1** click **Step 1: Time Dependent**.
- 2 In the **Settings** window for **Time Dependent**, locate the **Study Settings** section.
- 3 In the **Output times** text field, type range (0,5.0e-8,2.0e-5).  
For all of the studies in this example, a tight user-defined tolerance will be used to ensure particle kinetic energy is conserved.
- 4 From the **Tolerance** list, choose **User controlled**.
- 5 In the **Relative tolerance** text field, type 1.0E-6.
- 6 In the **Model Builder** window, click **Study 1**.
- 7 In the **Settings** window for **Study**, type Lagrangian Study in the **Label** text field.
- 8 In the **Study** toolbar, click  **Compute**.

## RESULTS

### *Lagrangian Results*

In order to be able to see the radius of the particle orbit, plot the  $y$ -component of the particle location as a color expression.

- 1 In the **Settings** window for **3D Plot Group**, type Lagrangian Results in the **Label** text field.
- 2 Locate the **Plot Settings** section. Clear the **Plot dataset edges** checkbox.



### *Particle Trajectories 1*

Render the particle trajectory as a ribbon. The default ribbon orientation is in the direction of the unit binormal, or the direction out of the plane tangent to the curved trajectory.


- 1 In the **Model Builder** window, expand the **Lagrangian Results** node, then click **Particle Trajectories 1**.
- 2 In the **Settings** window for **Particle Trajectories**, locate the **Coloring and Style** section.
- 3 Find the **Line style** subsection. From the **Type** list, choose **Ribbon**.
- 4 Select the **Width scale factor** checkbox. In the associated text field, type 4E-5.
- 5 Find the **Point style** subsection. From the **Type** list, choose **None**.

### *Color Expression 1*


- 1 In the **Model Builder** window, expand the **Particle Trajectories 1** node, then click **Color Expression 1**.

- 2 In the **Settings** window for **Color Expression**, locate the **Expression** section.
- 3 In the **Expression** text field, type  $qy/2$ .
- 4 Locate the **Coloring and Style** section. From the **Color table** list, choose **Inferno**.
- 5 In the **Lagrangian Results** toolbar, click  **Plot**.
- 6 Click the  **Zoom Extents** button in the **Graphics** toolbar. The plot should look like [Figure 1](#).

#### *Particle Evaluation 1*

- 1 In the **Results** toolbar, click  **More Derived Values** and choose **Other > Particle Evaluation**.
- 2 In the **Settings** window for **Particle Evaluation**, locate the **Data** section.
- 3 From the **Time selection** list, choose **First**.
- 4 Locate the **Expression** section. In the **Expression** text field, type  $rL$ .
- 5 Click the arrow next to the **Evaluate** button and click **New Table**.

#### *Particle Evaluation 2*

- 1 In the **Results** toolbar, click  **More Derived Values** and choose **Other > Particle Evaluation**.
- 2 In the **Settings** window for **Particle Evaluation**, locate the **Data** section.
- 3 From the **Time selection** list, choose **First**.
- 4 Locate the **Expression** section. In the **Expression** text field, type  $\text{timemax}(0, 2E-5, qy) / 2$ .
- 5 Click the arrow next to the **Evaluate** button and select **Table 1 - Particle Evaluation 1 (rL)**.

Now switch formulation from **Lagrangian** to **Hamiltonian**. When you do this, the particle momentum components are added as additional degrees of freedom. The momentum has three components:  $p_x$ ,  $p_y$ , and  $p_z$ . This results in a doubling of the number of degrees of freedom in the model.



### **MATHEMATICAL PARTICLE TRACING (PT)**

- 1 In the **Model Builder** window, under **Component 1 (comp1)** click **Mathematical Particle Tracing (pt)**.
- 2 In the **Settings** window for **Mathematical Particle Tracing**, locate the **Particle Release and Propagation** section.
- 3 From the **Formulation** list, choose **Hamiltonian**.

### Particle Properties 1


- 1 In the **Model Builder** window, under **Component 1 (comp1) > Mathematical Particle Tracing (pt)** click **Particle Properties 1**.
- 2 In the **Settings** window for **Particle Properties**, locate the **Hamiltonian** section.
- 3 In the  $H$  text field, type  $((px - e\_const * Ax)^2 + (py - e\_const * Ay)^2 + (pz - e\_const * Az)^2) / (2 * pt.mp)$ .

### ADD STUDY

- 1 In the **Home** toolbar, click  **Add Study** to open the **Add Study** window.
- 2 Go to the **Add Study** window.
- 3 Find the **Studies** subsection. In the **Select Study** tree, select **General Studies > Time Dependent**.
- 4 Click the **Add Study** button in the window toolbar.
- 5 In the **Home** toolbar, click  **Add Study** to close the **Add Study** window.

### STUDY 2

#### Step 1: Time Dependent

- 1 In the **Settings** window for **Time Dependent**, locate the **Study Settings** section.
- 2 In the **Output times** text field, type `range(0, 5.0e-8, 2.0e-5)`.
- 3 From the **Tolerance** list, choose **User controlled**.
- 4 In the **Relative tolerance** text field, type `1.0E-6`.
- 5 In the **Model Builder** window, click **Study 2**.
- 6 In the **Settings** window for **Study**, type `Hamiltonian Study` in the **Label** text field.
- 7 In the **Study** toolbar, click  **Compute**.

### RESULTS

#### Hamiltonian Results



- 1 In the **Settings** window for **3D Plot Group**, type `Hamiltonian Results` in the **Label** text field.
- 2 Locate the **Plot Settings** section. Clear the **Plot dataset edges** checkbox.

#### Particle Trajectories 1

- 1 In the **Model Builder** window, expand the **Hamiltonian Results** node, then click **Particle Trajectories 1**.
- 2 In the **Settings** window for **Particle Trajectories**, locate the **Coloring and Style** section.

- 3 Find the **Line style** subsection. From the **Type** list, choose **Ribbon**.
- 4 Select the **Width scale factor** checkbox. In the associated text field, type 4E-5.
- 5 Find the **Point style** subsection. From the **Type** list, choose **None**.

#### *Color Expression 1*

- 1 In the **Model Builder** window, expand the **Particle Trajectories 1** node, then click **Color Expression 1**.
- 2 In the **Settings** window for **Color Expression**, locate the **Expression** section.
- 3 In the **Expression** text field, type  $qy/2$ .
- 4 Locate the **Coloring and Style** section. From the **Color table** list, choose **Inferno**.
- 5 In the **Hamiltonian Results** toolbar, click  **Plot**.
- 6 Click the  **Go to Default View** button in the **Graphics** toolbar. The plot should look like [Figure 2](#).

#### *Particle Evaluation 3*

- 1 In the **Model Builder** window, expand the **Results > Datasets** node.
- 2 Right-click **Results > Derived Values** and choose **More Derived Values > Particle Evaluation**.
- 3 In the **Settings** window for **Particle Evaluation**, locate the **Data** section.
- 4 From the **Dataset** list, choose **Particle 2**.
- 5 From the **Time selection** list, choose **First**.
- 6 Locate the **Expression** section. In the **Expression** text field, type  $\text{timemax}(0, 2E-5, qy) / 2$ .
- 7 Click the arrow next to the **Evaluate** button and select **Table 1 - Particle Evaluation 1 (rL)**.

Switch to the **Newtonian** formulation and add the Lorentz force manually.

### **MATHEMATICAL PARTICLE TRACING (PT)**

- 1 In the **Model Builder** window, under **Component 1 (comp1)** click **Mathematical Particle Tracing (pt)**.
- 2 In the **Settings** window for **Mathematical Particle Tracing**, locate the **Particle Release and Propagation** section.
- 3 From the **Formulation** list, choose **Newtonian**.



#### *Force 1*

- 1 In the **Physics** toolbar, click  **Domains** and choose **Force**.
- 2 In the **Settings** window for **Force**, locate the **Domain Selection** section.

- From the **Selection** list, choose **All domains**.
- Locate the **Force** section. Specify the  $\mathbf{F}$  vector as


$e\_const*(Bz*pt.vy-By*pt.vz)$	x
$e\_const*(-Bz*pt.vx+Bx*pt.vz)$	y
$e\_const*(By*pt.vx-Bx*pt.vy)$	z

### ADD STUDY

- In the **Home** toolbar, click  **Add Study** to open the **Add Study** window.
- Go to the **Add Study** window.
- Find the **Studies** subsection. In the **Select Study** tree, select **General Studies > Time Dependent**.
- Click the **Add Study** button in the window toolbar.
- In the **Home** toolbar, click  **Add Study** to close the **Add Study** window.

### STUDY 3

#### Step 1: Time Dependent

- In the **Settings** window for **Time Dependent**, locate the **Study Settings** section.
- In the **Output times** text field, type range (0,5.0e-8,2.0e-5).
- From the **Tolerance** list, choose **User controlled**.
- In the **Relative tolerance** text field, type 1.0E-6.
- In the **Model Builder** window, click **Study 3**.
- In the **Settings** window for **Study**, type Newtonian Study in the **Label** text field.
- In the **Study** toolbar, click  **Compute**.

### RESULTS

#### Newtonian Results



- In the **Settings** window for **3D Plot Group**, type Newtonian Results in the **Label** text field.
- Locate the **Plot Settings** section. Clear the **Plot dataset edges** checkbox.

#### Particle Trajectories I

- In the **Model Builder** window, expand the **Newtonian Results** node, then click **Particle Trajectories I**.
- In the **Settings** window for **Particle Trajectories**, locate the **Coloring and Style** section.

- 3 Find the **Line style** subsection. From the **Type** list, choose **Ribbon**.
- 4 Select the **Width scale factor** checkbox. In the associated text field, type 4E-5.
- 5 Find the **Point style** subsection. From the **Type** list, choose **None**.

#### *Color Expression 1*

- 1 In the **Model Builder** window, expand the **Particle Trajectories 1** node, then click **Color Expression 1**.
- 2 In the **Settings** window for **Color Expression**, locate the **Expression** section.
- 3 In the **Expression** text field, type  $qy/2$ .
- 4 Locate the **Coloring and Style** section. From the **Color table** list, choose **Inferno**.
- 5 In the **Newtonian Results** toolbar, click  **Plot**.
- 6 Click the  **Go to Default View** button in the **Graphics** toolbar. The plot should look like [Figure 3](#).

#### *Particle Evaluation 4*


- 1 In the **Results** toolbar, click  $\frac{8.85}{e-12}$  **More Derived Values** and choose **Other > Particle Evaluation**.
- 2 In the **Settings** window for **Particle Evaluation**, locate the **Data** section.
- 3 From the **Dataset** list, choose **Particle 3**.
- 4 From the **Time selection** list, choose **First**.
- 5 Locate the **Expression** section. In the **Expression** text field, type  $\text{timemax}(0, 2E-5, qy) / 2$ .
- 6 Click the arrow next to the **Evaluate** button and select **Table 1 - Particle Evaluation 1 (rL)**.


Finally, switch to the **Newtonian, first order** formulation.

### **MATHEMATICAL PARTICLE TRACING (PT)**

- 1 In the **Model Builder** window, under **Component 1 (comp1)** click **Mathematical Particle Tracing (pt)**.
- 2 In the **Settings** window for **Mathematical Particle Tracing**, locate the **Particle Release and Propagation** section.
- 3 From the **Formulation** list, choose **Newtonian, first order**.


### **ADD STUDY**

- 1 In the **Home** toolbar, click  **Add Study** to open the **Add Study** window.
- 2 Go to the **Add Study** window.

- 3 Find the **Studies** subsection. In the **Select Study** tree, select **General Studies > Time Dependent**.
- 4 Click the **Add Study** button in the window toolbar.
- 5 In the **Home** toolbar, click  **Add Study** to close the **Add Study** window.

## STUDY 4

### *Step 1: Time Dependent*

- 1 In the **Settings** window for **Time Dependent**, locate the **Study Settings** section.
- 2 In the **Output times** text field, type range (0,5.0e-8,2.0e-5).
- 3 From the **Tolerance** list, choose **User controlled**.
- 4 In the **Relative tolerance** text field, type 1.0E-6.
- 5 In the **Model Builder** window, click **Study 4**.
- 6 In the **Settings** window for **Study**, type Newtonian, First Order Study in the **Label** text field.
- 7 In the **Study** toolbar, click  **Compute**.

## RESULTS

### *Newtonian, First Order Results*



- 1 In the **Settings** window for **3D Plot Group**, type Newtonian, First Order Results in the **Label** text field.
- 2 Locate the **Plot Settings** section. Clear the **Plot dataset edges** checkbox.

### *Particle Trajectories 1*


- 1 In the **Model Builder** window, expand the **Newtonian, First Order Results** node, then click **Particle Trajectories 1**.
- 2 In the **Settings** window for **Particle Trajectories**, locate the **Coloring and Style** section.
- 3 Find the **Line style** subsection. From the **Type** list, choose **Ribbon**.
- 4 Select the **Width scale factor** checkbox. In the associated text field, type 4E-5.
- 5 Find the **Point style** subsection. From the **Type** list, choose **None**.

### *Color Expression 1*

- 1 In the **Model Builder** window, expand the **Particle Trajectories 1** node, then click **Color Expression 1**.
- 2 In the **Settings** window for **Color Expression**, locate the **Expression** section.
- 3 In the **Expression** text field, type  $qy/2$ .

- 4 Locate the **Coloring and Style** section. From the **Color table** list, choose **Inferno**.
- 5 In the **Newtonian, First Order Results** toolbar, click  **Plot**.
- 6 Click the  **Go to Default View** button in the **Graphics** toolbar. The plot should look like [Figure 4](#).

#### *Particle Evaluation 5*

- 1 In the **Results** toolbar, click  **More Derived Values** and choose **Other > Particle Evaluation**.
- 2 In the **Settings** window for **Particle Evaluation**, locate the **Data** section.
- 3 From the **Dataset** list, choose **Particle 4**.
- 4 From the **Time selection** list, choose **First**.
- 5 Locate the **Expression** section. In the **Expression** text field, type  $\text{timemax}(0, 2E-5, \text{qy}) / 2$ .
- 6 Click the arrow next to the **Evaluate** button and select **Table 1 - Particle Evaluation 1 (rL)**.  
The Larmor radius shows good agreement with the analytic expression for all four formulations.