



Model created in COMSOL Multiphysics 6.4

# Buoyancy Flow with Darcy's Law – The Elder Problem

## Introduction

---

Density variations can initiate flow even in a still fluid. In earth systems, density variations can arise from naturally occurring salts, subsurface temperature changes, or migrating pollution. This buoyant or density-driven flow factors into fluid movement in salt-lake systems, saline-disposal basins, dense contaminant and leachate plumes, and geothermal reservoirs, to name just a few.


This example duplicates a benchmark problem for time-dependent buoyant flow in a porous medium. Known as the Elder problem (Ref. 1), it follows a laboratory experiment to study thermal convection. When Voss and Souza (Ref. 2) recast the Elder problem for salt concentrations, it became a benchmark that many researchers (Ref. 8) have used to test a number of variable-density flow codes including SUTRA (Ref. 3) and SEAWAT (Ref. 4).

This application examines the Elder problem for concentrations by coupling two physics interfaces: Darcy's Law and Transport of Diluted Species in Porous Media.

## Model Definition

---

In this example (Figure 1) a vertical cross section of a water-saturated porous medium extends 300 m in the  $x$  direction and 150 m in the  $y$  direction.



*Figure 1: Geometry for modeling the Elder problem with initial conditions and boundary conditions indicated. In this cross section of a water-saturated porous medium, high salt concentrations exist in the top-right region.*

The material properties are homogeneous and isotropic. A vertical line at  $x = 300$  m represents a symmetry boundary with a mirror image of the cross section extending

beyond it. There is no flow across the geometry edges. High salt concentrations exist at the upper boundary (along  $y = 150$  m) from  $x = 150$  m to 300 m. Salt concentration is zero along the lower boundary. The water is initially stationary (with a hydrostatic pressure distribution) and pristine (zero salt concentration). When the density increases near the high-concentration boundary, flow develops. The period of interest is 20 years. According to Ref. 6 and Ref. 7, the length chosen by Elder is close to a critical value that separates downwelling and upwelling plume structures. As a consequence, the problem is particularly sensitive to perturbations.

### FLUID FLOW

You can describe the fluid flow in this problem using Darcy's law:

$$\frac{\partial \epsilon_p \rho}{\partial t} + \nabla \cdot \rho \mathbf{u} = 0$$

$$\mathbf{u} = -\frac{\kappa}{\mu} (\nabla p + \rho g \nabla D)$$

Here,  $\rho$  is the water density ( $\text{kg}/\text{m}^3$ ),  $t$  is the time (s),  $\epsilon_p$  is the porosity, and  $\mathbf{u}$  is the vector of directional seepage rates, also known as the Darcy velocity. The Darcy velocity  $\mathbf{u}$  depends on the permeability  $\kappa$  ( $\text{m}^2$ ), the fluid's dynamic viscosity  $\mu$  (Pa·s), the fluid's pressure  $p$  (Pa), and the acceleration of gravity  $g$  ( $\text{m}/\text{s}^2$ ). The gradient of the elevation  $D$  (m) indicates the direction of the vertical coordinate,  $y$ .

In Elder's problem, the fluid density depends linearly on the salt concentration,  $c$  ( $\text{mol}/\text{m}^3$ ) according to

$$\rho = \rho_0 + \beta c = \rho_0 + \frac{\rho_s - \rho_0}{c_s - c_0} c$$

Here,  $c_0$  and  $c_s$  are the normalized salt concentrations of pristine and salty water.

The symmetry or zero flow on all boundaries fix only the change in pressure. For a unique solution, you must also specify a reference pressure. In this example, the pressure at the point (0,0) is fixed. With the Darcy's Law interface, you express all these conditions as

$$\begin{aligned} \mathbf{n} \cdot \rho \mathbf{u} &= 0 & \partial\Omega & \text{ Sides} \\ p &= p_0 & \partial^2\Omega & \text{ Point} \\ p(x, y, 0) &= \rho_0 g D & t = 0 & \text{ Initial value} \end{aligned}$$

where  $\mathbf{n}$  is the unit vector normal to the boundary, and  $p_0$  is the reference pressure.

## TRANSPORT OF DILUTED SPECIES IN POROUS MEDIA

The governing equation for this problem is the conservative form of the Transport of Diluted Species in Porous Media interface

$$\frac{\partial \theta_s c}{\partial t} + \mathbf{u} \cdot \nabla c - \nabla \cdot \theta_s \tau D_L \nabla c = 0$$

where  $D_L$  is the fluid's diffusion coefficient ( $\text{m}^2/\text{d}$ );  $\theta_s$  is the fluid's volume fraction (porosity);  $c$  is the salt concentration ( $\text{mol}/\text{m}^3$ ), and  $\mathbf{u}$  is the Darcy velocity ( $\text{m}/\text{s}$ ).

In Elder's problem, the contaminant spreads only by advection and molecular diffusion, and the salt concentration is normalized to unit values.

The only contaminant source in the model domain is the salt concentration along the right half of the upper boundary. The vertical edge at  $x = 300$  m is a symmetry boundary. The remaining boundaries have zero flux. The initial concentration is zero. The following equations represent these conditions:

$$\begin{aligned} \mathbf{n} \cdot [c\mathbf{u} - \theta_s \tau D_L \nabla c] &= 0 & \partial\Omega \text{ Sides} \\ c &= c_s & \partial\Omega \text{ Salt} \\ c(x, y, 0) &= 0 & t = 0 \text{ Initial value} \end{aligned}$$

where  $\mathbf{n}$  is the unit vector normal to the boundary.

### *Model Data*

The example works with the following data:

PARAMETER	NAME	VALUE
$\rho_0$	Fresh-water density	1000 $\text{kg}/\text{m}^3$
$\rho_s$	Salt-water density	1200 $\text{kg}/\text{m}^3$
$\kappa$	Permeability	0.5 Darcy
$\mu$	Dynamic viscosity	0.001 Pa·s
$g$	Gravity	9.81 $\text{m}/\text{s}^2$
$\varepsilon$	Porosity	0.1
$\tau_L D_L$	Molecular diffusion rate	$3.56 \cdot 10^{-6} \text{ m}^2/\text{s}$
$c_s$	Salt-water concentration	1 $\text{mol}/\text{m}^3$

PARAMETER	NAME	VALUE
$c_0$	Fresh-water concentration	0 mol/m <sup>3</sup>
$\beta$	Increase of water density due to concentration changes	200 kg/mol

Note that the original set of parameters for the Elder problem gives a global Peclet number equal to

$$\text{Pe} = \frac{(\rho_s - \rho_0)g\kappa L}{\mu\varepsilon_p D_L} \sim 408$$


Here, the difference in water density is  $\rho_s - \rho_0 = \beta(c_s - c_0) = 200 \text{ kg/m}^3$ , and the length scale  $L$  is 150 m. This high Peclet number poses extra difficulties to numerical schemes; see for instance [Ref. 5](#) to [Ref. 7](#).

### *Results and Discussion*

---

The following results come from the COMSOL Multiphysics solution to a benchmark buoyancy problem that is often used for both temperatures ([Ref. 1](#)) and concentrations ([Ref. 2](#)).

[Figure 2](#) gives snapshots of concentrations at six times during the 20-year simulation period. Initially the water is pristine. By the end of the first year, concentrations spread by diffusion, creating a density gradient. The buoyancy flow begins at the edge of the salt contact, where there is a sharp contrast in fluid density. By the end of year three, the fingering of high concentrations into the reservoir is mature. By year 10, the salt concentrations have spread over roughly 60% of the model domain. The COMSOL Multiphysics solution in [Figure 2](#) is in excellent agreement with that from Elder ([Ref. 1](#)).



*Figure 2: Snapshots of concentrations from the COMSOL Multiphysics solution to the buoyancy-flow benchmark developed by Voss and Souza (Ref. 2) for the Elder problem.*

Figure 3 shows the concentration for the stationary Elder’s problem.





Figure 3: Concentration distribution from the COMSOL Multiphysics solution to the buoyancy-flow benchmark for the stationary Elder problem.

Of interest in the Elder problem is the development of convection cells. The COMSOL Multiphysics plots in Figure 4 reveal the convection cells with the help of velocity streamlines, which the figure shows simultaneously with concentrations for years 3, 10, 15,

and 20. At early times, small convection cells develop between the individual fingers of the plume. At late times, a single convection cell covers the model domain.



*Figure 4: Salt concentrations (surface plot) and velocities (streamlines) from the COMSOL Multiphysics solution to a buoyancy benchmark problem (Ref. 2).*

This example shows COMSOL Multiphysics applied to a well-known benchmark problem applicable to flow driven by density variations related to either temperature or concentration. The COMSOL Multiphysics results, here for concentration, closely match the benchmark solution (Ref. 2). This buoyant flow is straightforward to set up directly on top of a standard fluid flow and solute-transport model.

## *References*

1. J.W. Elder, “Transient Convection in a Porous Medium,” *J. Fluid Mech.*, vol. 27, no. 3, pp. 609–623, 1967.
2. C.I. Voss and W.R. Souza, “Variable Density Flow and Solute Transport Simulation of Regional Aquifers Containing a Narrow Freshwater-saltwater Transition Zone,” *Water Resour. Res.*, vol. 23, no. 10, pp. 1851–1866, 1987.
3. C.I. Voss, “A Finite-element Simulation Model for Saturated-unsaturated, Fluid-density-dependent Ground-water Flow with Energy Transport or Chemically-reactive Single-species Solute Transport,” *U.S. Geological Survey Water-Resources Investigation Report*, 84-4369, 1984.

4. W. Guo and C.D. Langevin, *User's Guide to SEAWAT: A Computer Program for Simulation of Three-Dimensional Variable-Density Ground-Water Flow*, U.S. Geological Survey Techniques of Water-Resources Investigations 6-A7, 2002.
5. P. Frolkovic and H. De Schepper, "Numerical modelling of convection dominated transport coupled with density driven flow in porous media," *Adv. Water Resour.*, vol. 24, no. 1, pp. 63–72, 2000.
6. G.F. Carey and others, "Modelling error and constitutive relations in simulation of flow and transport," *Int. J. Numer. Meth. Fluids*, vol. 46, pp. 1211–1236, 2004.
7. J.A. Woods and G.F. Carey, "Upwelling and downwelling behavior in the Elder-Voss-Souza benchmark," *Water Resour. Res.*, vol. 43, W12403, 2007.
8. J.W. Elder and others, "The Elder Problem," *Fluids*, vol. 2, p. 11, 2017.

---

**Application Library path:** Subsurface\_Flow\_Module/Solute\_Transport/  
buoyancy\_darcy\_elder


---

### *Modeling Instructions*




---

From the **File** menu, choose **New**.

#### **NEW**

In the **New** window, click  **Model Wizard**.


#### **MODEL WIZARD**

- 1 In the **Model Wizard** window, click  **2D**.
- 2 In the **Select Physics** tree, select **Fluid Flow > Porous Media and Subsurface Flow > Darcy's Law (dl)**.
- 3 Click **Add**.
- 4 In the **Select Physics** tree, select **Chemical Species Transport > Transport of Diluted Species in Porous Media (tds)**.
- 5 Click **Add**.
- 6 Click  **Study**.
- 7 In the **Select Study** tree, select **General Studies > Time Dependent**.
- 8 Click  **Done**.

## GLOBAL DEFINITIONS


Start by loading parameters from a file.

### *Parameters 1*



- 1 In the **Model Builder** window, under **Global Definitions** click **Parameters 1**.
- 2 In the **Settings** window for **Parameters**, locate the **Parameters** section.
- 3 Click  **Load from File**.
- 4 Browse to the model's Application Libraries folder and double-click the file `buoyancy_darcy_elder_parameters.txt`.

## GEOMETRY 1

### *Rectangle 1 (r1)*

- 1 In the **Geometry** toolbar, click  **Rectangle**.
- 2 In the **Settings** window for **Rectangle**, locate the **Size and Shape** section.
- 3 In the **Width** text field, type  $2*L$ .
- 4 In the **Height** text field, type  $L$ .


### *Point 1 (pt1)*

- 1 In the **Geometry** toolbar, click  **Point**.
- 2 In the **Settings** window for **Point**, locate the **Point** section.
- 3 In the **x** text field, type  $L$ .
- 4 In the **y** text field, type  $L$ .
- 5 Click  **Build All Objects**.

## DEFINITIONS

Add a variable for the buoyancy force due to concentration gradients.

### *Variables 1*

- 1 In the **Definitions** toolbar, click  **Local Variables**.
- 2 In the **Settings** window for **Variables**, locate the **Variables** section.
- 3 In the table, enter the following settings:

Name	Expression	Unit	Description
rho	$\text{rho}0 + \text{beta} * c * (c > 0)$	kg/m <sup>3</sup>	Water density

## DARCY'S LAW (DL)

- 1 In the **Model Builder** window, under **Component 1 (comp1)** click **Darcy's Law (dl)**.

- 2 In the **Settings** window for **Darcy's Law**, locate the **Gravity Effects** section.
- 3 Select the **Include gravity** checkbox.

#### *Gravity I*

- 1 In the **Model Builder** window, under **Component 1 (comp1) > Darcy's Law (dl)** click **Gravity I**.
- 2 In the **Settings** window for **Gravity**, locate the **Gravity** section.
- 3 From the **Specify** list, choose **Elevation**.
- 4 Select the **Specify reference position** checkbox.
- 5 Specify the  $\mathbf{r}_{\text{ref}}$  vector as

0	x
L	y

#### *Fluid I*

- 1 In the **Model Builder** window, under **Component 1 (comp1) > Darcy's Law (dl) > Porous Medium I** click **Fluid I**.
- 2 In the **Settings** window for **Fluid**, locate the **Fluid Properties** section.
- 3 From the  $\rho$  list, choose **User defined**. In the associated text field, type  $\rho$ .
- 4 From the  $\mu$  list, choose **User defined**. In the associated text field, type  $\mu$ .

#### *Porous Matrix I*

- 1 In the **Model Builder** window, click **Porous Matrix I**.
- 2 In the **Settings** window for **Porous Matrix**, locate the **Matrix Properties** section.
- 3 From the  $\epsilon_p$  list, choose **User defined**. In the associated text field, type  $\epsilon$ .
- 4 From the  $\kappa$  list, choose **User defined**. In the associated text field, type  $\kappa$ .

#### *Initial Values I*

- 1 In the **Model Builder** window, under **Component 1 (comp1) > Darcy's Law (dl)** click **Initial Values I**.
- 2 In the **Settings** window for **Initial Values**, locate the **Initial Values** section.
- 3 Click the **Hydraulic head** button.


With gravity active, an initial zero hydraulic head defines the hydrostatic pressure distribution as reasonable initial condition.

#### *Symmetry I*

- 1 In the **Physics** toolbar, click  **Boundaries** and choose **Symmetry**.

2 Select Boundary 5 only.

Since the pressure is not set explicitly by a boundary condition, you need to fix it at least at one point to get a unique solution for Darcy's Law.

3 Click the  **Show More Options** button in the **Model Builder** toolbar.

4 In the **Show More Options** dialog, in the tree, select the checkbox for the node **Physics > Equation Contributions**.

5 Click **OK**.

#### *Pointwise Constraint 1*

1 In the **Physics** toolbar, click  **Points** and choose **Pointwise Constraint**.

2 In the **Settings** window for **Pointwise Constraint**, locate the **Pointwise Constraint** section.

3 In the **Constraint expression** text field, type  $p_0 - p$ .

4 Select Point 2 only.

### **TRANSPORT OF DILUTED SPECIES IN POROUS MEDIA (TDS)**

Continue with setting up the species transport interface.

#### *Fluid 1*

1 In the **Model Builder** window, under **Component 1 (comp1) > Transport of Diluted Species in Porous Media (tds) > Porous Medium 1** click **Fluid 1**.

2 In the **Settings** window for **Fluid**, locate the **Convection** section.

3 From the **u** list, choose **Total Darcy velocity field (dl/porous1)**.

4 Locate the **Diffusion** section. In the  $D_{F,c}$  text field, type  $D_L$ .

5 From the **Effective diffusivity model** list, choose **Tortuosity model**.

#### *Porous Matrix 1*

1 In the **Model Builder** window, click **Porous Matrix 1**.

2 In the **Settings** window for **Porous Matrix**, locate the **Matrix Properties** section.

3 From the  $\epsilon_p$  list, choose **User defined**. In the associated text field, type  $\epsilon$ .

#### *Initial Values 1*

1 In the **Model Builder** window, under **Component 1 (comp1) > Transport of Diluted Species in Porous Media (tds)** click **Initial Values 1**.


2 In the **Settings** window for **Initial Values**, locate the **Initial Values** section.

3 In the  $c$  text field, type  $c_0$ .


### *Symmetry 1*

- 1 In the **Physics** toolbar, click  **Boundaries** and choose **Symmetry**.
- 2 Select Boundary 5 only.

### *Concentration 1*


- 1 In the **Physics** toolbar, click  **Boundaries** and choose **Concentration**.
- 2 Select Boundary 2 only.
- 3 In the **Settings** window for **Concentration**, locate the **Concentration** section.
- 4 Select the **Species c** checkbox.
- 5 In the  $c_{0,c}$  text field, type  $c_0$ .

### *Concentration 2*


- 1 In the **Physics** toolbar, click  **Boundaries** and choose **Concentration**.
- 2 Select Boundary 4 only.
- 3 In the **Settings** window for **Concentration**, locate the **Concentration** section.
- 4 Select the **Species c** checkbox.
- 5 In the  $c_{0,c}$  text field, type  $c_s$ .

## **MESH 1**

### *Mapped 1*

In the **Mesh** toolbar, click  **Mapped**.

### *Size*

- 1 In the **Model Builder** window, click **Size**.
- 2 In the **Settings** window for **Size**, locate the **Element Size** section.
- 3 From the **Predefined** list, choose **Extremely fine**.
- 4 Click  **Build All**.



## **STUDY 1**

### *Step 1: Time Dependent*

- 1 In the **Model Builder** window, under **Study 1** click **Step 1: Time Dependent**.
- 2 In the **Settings** window for **Time Dependent**, locate the **Study Settings** section.
- 3 From the **Time unit** list, choose **a**.
- 4 In the **Output times** text field, type  $\text{range}(0, 1, 20)$ .

It is a good idea to restrict the maximum time step size to capture the convective motion accurately.

#### *Solution 1 (sol1)*

- 1 In the **Study** toolbar, click  **Show Default Solver**.
- 2 In the **Model Builder** window, expand the **Solution 1 (sol1)** node, then click **Time-Dependent Solver 1**.
- 3 In the **Settings** window for **Time-Dependent Solver**, click to expand the **Time Stepping** section.
- 4 From the **Maximum step constraint** list, choose **Constant**.
- 5 Click  **Run**.

## **RESULTS**

#### *Pressure (dl)*

The first default plot group shows the pressure distribution, the second the velocity field.

#### *Concentration (tds)*

The third default plot group shows the concentration after 20 years. To reproduce the series of plots in [Figure 2](#), add contours and plot for different times.

#### *Contour 1*

- 1 In the **Model Builder** window, right-click **Concentration (tds)** and choose **Contour**.
- 2 In the **Settings** window for **Contour**, click **Replace Expression** in the upper-right corner of the **Expression** section. From the menu, choose **Component 1 (comp1) > Transport of Diluted Species in Porous Media > Species c > c - Molar concentration, c - mol/m<sup>3</sup>**.
- 3 Locate the **Coloring and Style** section. Clear the **Color legend** checkbox.
- 4 From the **Coloring** list, choose **Uniform**.
- 5 From the **Color** list, choose **Black**.

#### *Arrow Surface 1*

In the **Model Builder** window, right-click **Arrow Surface 1** and choose **Disable**.

#### *Concentration (tds)*

- 1 In the **Model Builder** window, click **Concentration (tds)**.
- 2 In the **Settings** window for **2D Plot Group**, click to expand the **Title** section.
- 3 From the **Title type** list, choose **Manual**.
- 4 In the **Title** text area, type Concentration (kg/m<sup>3</sup>).

5 Locate the **Data** section. From the **Time (a)** list, choose **1**.

6 In the **Concentration (tds)** toolbar, click  **Plot**.

Compare the result with the upper-left plot in [Figure 2](#).

Repeat the previous two steps for 2 years, 3 years, 10 years, 15 years, and 20 years to reproduce the remaining five plots in the series.

To reproduce the combined concentration/velocity plots in [Figure 4](#), proceed as follows.

#### *Concentration (tds) I*

Right-click **Concentration (tds)** and choose **Duplicate**.

#### *Contour I*

1 In the **Model Builder** window, expand the **Concentration (tds) I** node.


2 Right-click **Contour I** and choose **Delete**.

3 Click **Yes** to confirm.

#### *Surface I*

In the **Model Builder** window, under **Results > Concentration (tds) I** click **Surface I**.

#### *Concentration (tds) I*

In the **Concentration (tds) I** toolbar, click  **Streamline**.

#### *Streamline I*

1 In the **Settings** window for **Streamline**, locate the **Streamline Positioning** section.

2 From the **Positioning** list, choose **Uniform density**.

3 Locate the **Coloring and Style** section. Find the **Point style** subsection. From the **Type** list, choose **Arrow**.

4 From the **Arrow length** list, choose **Logarithmic**.

5 From the **Color** list, choose **Gray**.

#### *Concentration and Velocity*

1 In the **Model Builder** window, under **Results** click **Concentration (tds) I**.

2 In the **Settings** window for **2D Plot Group**, type **Concentration** and **Velocity** in the **Label** text field.

3 Locate the **Data** section. From the **Time (a)** list, choose **3**.

4 Locate the **Title** section. In the **Title** text area, type **Surface: Concentration (kg/m<sup>3</sup>) Streamlines: Velocity field**.

5 In the **Concentration and Velocity** toolbar, click  **Plot**.

Compare the result with the upper-left plot in [Figure 4](#).

Repeat the previous two steps for 10 years, 15 years, and 20 years to reproduce the remaining three plots in the series.

## ROOT

Finally compute the stationary solution to the Elder problem.

## ADD STUDY

1 In the **Home** toolbar, click  **Add Study** to open the **Add Study** window.

2 Go to the **Add Study** window.

3 Find the **Studies** subsection. In the **Select Study** tree, select **General Studies > Stationary**.

4 Click the **Add Study** button in the window toolbar.

5 In the **Home** toolbar, click  **Add Study** to close the **Add Study** window.

## STUDY 2

1 In the **Settings** window for **Study**, locate the **Study Settings** section.

2 Clear the **Generate default plots** checkbox.

### *Step 1: Stationary*

1 In the **Model Builder** window, under **Study 2** click **Step 1: Stationary**.


2 In the **Settings** window for **Stationary**, click to expand the **Values of Dependent Variables** section.

3 Find the **Initial values of variables solved for** subsection. From the **Settings** list, choose **User controlled**.

4 From the **Method** list, choose **Solution**.

5 From the **Study** list, choose **Study 1, Time Dependent**.

The solution at the last time step is a good starting point for computing the stationary solution.


6 In the **Study** toolbar, click  **Compute**.

## RESULTS

To visualize the stationary concentration distribution, use the **Concentration** plot as starting point.

### *Concentration, Stationary*

1 In the **Model Builder** window, right-click **Concentration (tds)** and choose **Duplicate**.

- 2 In the **Settings** window for **2D Plot Group**, type Concentration, Stationary in the **Label** text field.
- 3 Locate the **Data** section. From the **Dataset** list, choose **Study 2/Solution 2 (sol2)**.
- 4 In the **Concentration, Stationary** toolbar, click  **Plot**.  
Compare the resulting plot with that in [Figure 3](#).